

Serosurvey of Roe Deer, Chamois and Domestic Sheep in the Central Italian Alps

Alessandra Gaffuri,^{1,6} Marco Giacometti,² Vito Massimo Tranquillo,³ Simone Magnino,⁴ Paolo Cordioli,³ and Paolo Lanfranchi⁵ ¹ Istituto Zooprofilattico Sperimentale della Lombardia e dell'Emilia Romagna "Bruno Ubertini" (IZSLER), Via Rovelli 53, I-24100 Bergamo, Italy; ² Wildvet Projects, CH-7605 Stampa, Switzerland; ³ IZSLER, Via Bianchi 9, I-25124 Brescia, Italy; ⁴ IZSLER, Via Taramelli 7, I-27100 Pavia, Italy; ⁵ Dipartimento di Patologia Animale, Igiene e Sanità Pubblica Veterinaria, Università di Milano, Via Celoria 10, I-20133 Milan, Italy; ⁶ Corresponding author (email: alessandra.gaffuri@bs.izs.it)

ABSTRACT: Roe deer (*Capreolus capreolus*), chamois (*Rupicapra rupicapra rupicapra*), and domestic sheep in the Orobie Alps, Italy, were serologically tested for antibodies to selected pathogens that may be transmitted across species. Antibodies against *Brucella* spp. and bovine herpesvirus 1 (roe deer and chamois only) were not detected in any species. In roe deer, antibodies were detected against *Toxoplasma gondii* (13%) and *Neospora caninum* (3%). Chamois tested positive for antibodies to *T. gondii* (5%), *N. caninum* (21%), bovine respiratory syncytial virus (BRSV) (41%), bovine parainfluenza type-3 virus (17%), pestiviruses (18%), and *Mycoplasma conjunctivae* (17%). In the sheep, particularly high antibody prevalence rates were found for *T. gondii* (78%), *Chlamydomphila* spp. (20%), pestiviruses (90%), BRSV (82%), and *M. conjunctivae* (81%).

Key words: Alpine chamois, domestic sheep, Italian Alps, roe deer, serosurvey.

Interactions between domestic and wild ungulates represent a potential problem in the Alps, due to livestock summer pasturing. Infectious diseases such as brucellosis (Ferroglio et al., 2000) and infectious keratoconjunctivitis (IKC; Belloy et al., 2003) have been found in wild ruminants due to transmission from livestock. In the Alps, sheep are of particular concern due to large herd sizes and the broad movements that often are associated with sheep management. To better understand the potential for pathogen transmission between sheep and wild ungulates in the Alps, a cross-sectional serologic survey of roe deer (*Capreolus capreolus*), chamois (*Rupicapra rupicapra rupicapra*), and sheep was done during 1998–2001. The study area (45°40' to 46°10'N, 9°25' to 10°20'E), situated in the Orobie Alps

(northern Italy), has three main valleys (Val Brembana, Val Seriana, and Val di Scalve). There are approximately 4,000 roe deer and 5,000 chamois in this area with the highest densities in Val Brembana; an outbreak of pneumonia and IKC recently caused high mortality in the chamois population in Val Brembana. Large wandering sheep flocks as well as smaller groups (approximately 25,000 sheep in total) share habitats with wild ungulates from June to September.

Blood samples were collected by hunters from 207 roe deer and 236 chamois harvested during the 1998–2001 hunting seasons. Refrigerated blood samples were delivered to the laboratory within 2 days; they were centrifuged and sera were stored at –20 C. Sera were collected from 13,565 sheep in 2000 and were stored at –20 C. From these samples, 352 sheep sera from each valley were randomly selected using a computerized random list produced with the "Random" procedure of the PEPI 3.01 software (Abramson and Gahlinger, 1993–2000). For sample size determination (sheep), we used an expected prevalence of 50% for all the tested pathogens; and based on Graat et al. (1997), this sample size provides a precision of 5% at a 95% confidence level.

Wildlife species were tested for antibodies to *Brucella* spp. by using complement fixation test (CFT) (Ciuchini and Farina, 1991) and the Rose Bengal agglutination test (D.M. 2 Luglio 1992 n.453) depending on the quality and amount of the sera. Ovine sera were tested by CFT; antibody titers ≥ 20 were considered positive.

For *Toxoplasma gondii* antibody testing, a commercial latex agglutination test (Toxotest-MT[®], Eiken Chemical Co., Tokyo, Japan) was used with a cut-off titer of 32 (Walls and Remington, 1983). Antibodies to *Neospora caninum* in wildlife species were detected with the indirect fluorescent antibody test performed on antigen slides (Conrad et al., 1993) with species-specific conjugates. Antigen consisted of tachyzoites of the NC-PV1 (*N. caninum*) Italian isolate (Magnino et al., 1999). Antibody titers were determined as the last dilution showing whole parasite fluorescence. For sheep, a commercial enzyme-linked immunoadsorbent assay (ELISA; Chekit Neospora[®], Bommeli Diagnostics, Liebefeld-Bern, Switzerland) was used.

Roe deer and chamois were tested for *Chlamydophila* spp. antibodies by using a CFT (O.I.E. Manual, 1992); titers ≥ 16 were considered positive. Sheep sera were tested using a commercial ELISA (Chekit Chlamydia[®], Bommeli Diagnostics).

A commercial blocking ELISA (Herd-Check Anti-IBR gE[®], IDEXX laboratories, Westbrook, Maine, USA) was used to test for antibodies to bovine herpesvirus 1 (BHV-1). Hemagglutination inhibition, by using SF-4 (ATCC VR281) as a reference virus, was used for bovine parainfluenza 3 virus (BPIV-3) antibody testing; 0.5% guinea pig red blood cells were used in this assay (Contini and Cossu, 1975). For pestivirus antibody testing, a blocking ELISA was performed using anti-p80 monoclonal antibodies as described previously (Brocchi et al., 1993); sera were tested at a 1:4 dilution. A blocking ELISA also was used to test for antibodies to bovine respiratory syncytial virus (BRSV; De Simone et al., 1986).

A commercial ELISA (Chekit *M. conjunctivae*[®], Bommeli Diagnostics) was used to test sheep for antibodies to *Mycoplasma conjunctivae* (Belloy et al., 2001). For chamois, a monoclonal antibody to sheep and goat IgG that was conjugated to horseradish peroxidase was used (Giacometti et al., 2002).

Because of inadequate serum quantity and poor quality due to hemolysis, individual sera were often not included in all assays. In particular, tests for antibodies against BHV-1 and BPIV-3 virus were not performed for the sheep.

For all antibody prevalence estimates, a 95% confidence interval (CI) was estimated according to a binomial distribution (Henken et al., 1997). Chi-square tests were used to test for differences in seroprevalence between valleys. The Cramer Φ coefficient was calculated according to Jekel et al. (1996) to value the effect of the sample size on the estimate of χ^2 .

Results are shown in Table 1. Negative results for *Brucella* spp. antibodies are consistent with observed absence of brucellosis in sheep in this area since 1997. In sheep, antibody prevalence was $>70\%$ for *T. gondii*, pestiviruses, respiratory syncytial virus, and *M. conjunctivae*. Significant differences in antibody prevalence between sheep sampled in different valleys were detected for *T. gondii* ($\chi^2=18.9_{(df=2)}$, $P<0.0001$), *Chlamydophila* spp. ($\chi^2=10.2_{(df=2)}$, $P=0.006$), and *M. conjunctivae* ($\chi^2=21.0_{(df=2)}$, $P<0.0001$; Table 2). In all cases, however, contingency coefficients were low (Cramer $\Phi_{\text{Toxoplasmosis}}=0.13$, Cramer $\Phi_{\text{Chlamydiosis}}=0.10$, and Cramer $\Phi_{\text{Keratoconjunctivitis}}=0.14$).

In roe deer, antibodies were only detected for *T. gondii* (13%) and *N. caninum* (3%); chamois also tested seropositive for *T. gondii* (5%) and *N. caninum* (21%). These protozoa infect a wide range of intermediate hosts, and definitive hosts, which include domestic cats, dogs, and other carnivores (Dubey, 1993; Gondim et al., 2004); these species are present in large parts of the study area. Our data agree with previously reported seropositive results from wild ruminants from the Alps (Gennero et al., 1993; Ferroglio and Rossi, 2001).

The prevalence of *M. conjunctivae*-seropositive chamois observed in this study was higher than prevalence estimates from the eastern Swiss Alps (Giacometti et al., 2002).

TABLE 1. Number of animals tested, positive animals, and overall seroprevalences for selected pathogens in three ruminants species in 1998–2001 in the Province of Bergamo (Italy).

| Pathogen | Roe deer | | | Chamois | | | Sheep | | |
|---------------------------|----------------|--------|-----------------------|----------------|--------|-----------------------|-----------------|--------|-----------------------|
| | No. of animals | | Prevalence (%) | No. of animals | | Prevalence (%) | No. of animals | | Prevalence (%) |
| | Positive | Tested | (95% CI) ^a | Positive | Tested | (95% CI) ^a | Positive | Tested | (95% CI) ^a |
| <i>Brucella</i> spp. | 0 | 50 | 0 (0.0–7.1) | 0 | 49 | 0 (0.0–7.2) | 0 | 1,056 | 0 (0.0–0.3) |
| <i>Toxoplasma gondii</i> | 27 | 207 | 13 (8.7–18.4) | 5 | 108 | 5 (1.5–10.4) | 821 | 1,056 | 78 (75.1–80.2) |
| <i>Neospora caninum</i> | 4 | 117 | 3 (0.9–8.5) | 14 | 67 | 21 (11.9–32.5) | 22 | 1,010 | 2 (1.3–3.2) |
| <i>Chlamydophila</i> spp. | 0 | 90 | 0 (0.0–4.0) | 0 | 106 | 0 (0.0–3.4) | 199 | 1,013 | 19.6 (17.2–22.2) |
| BHV-1 | 0 | 128 | 0 (0.0–2.8) | 0 | 100 | 0 (0.0–3.6) | nd ^b | nd | nd |
| BPIV-3 | 0 | 128 | 0 (0.0–2.8) | 15 | 90 | 17 (9.6–26.0) | nd | nd | nd |
| Pestiviruses | 0 | 138 | 0 (0.0–2.6) | 18 | 98 | 18 (11.2–27.4) | 954 | 1,056 | 90 (88.4–92.0) |
| BRSV | 0 | 131 | 0 (0.0–2.7) | 36 | 88 | 41 (30.5–51.9) | 879 | 1,056 | 82 (80.8–85.4) |
| <i>M. conjunctivae</i> | nd | nd | nd | 40 | 236 | 17 (12.5–22.4) | 857 | 1,056 | 81 (78.6–83.4) |

^a Binomial exact one-sided, 97.5% CI.^b nd=not done.

TABLE 2. Number of animals tested, positive animals, and seroprevalences for selected pathogens in sheep in 2000 in three valleys of the Province of Bergamo (Italy).

| Pathogen | Val Brembana | | | Val Seriana | | | Val di Scalve | | |
|---------------------------|----------------|--------|-----------------------|----------------|--------|-----------------------|----------------|--------|-----------------------|
| | No. of animals | | Prevalence (%) | No. of animals | | Prevalence (%) | No. of animals | | Prevalence (%) |
| | Positive | Tested | (95% CI) ^a | Positive | Tested | (95% CI) ^a | Positive | Tested | (95% CI) ^a |
| <i>Brucella</i> spp. | 0 | 352 | 0 (0.0–1.0) | 0 | 352 | 0 (0.0–1.0) | 0 | 352 | 0 (0.0–1.0) |
| <i>Toxoplasma gondii</i> | 301 | 352 | 86 (81.4–89.0) | 264 | 352 | 75 (70.1–79.4) | 256 | 352 | 73 (67.7–77.3) |
| <i>Neospora caninum</i> | 12 | 352 | 3 (1.7–5.8) | 6 | 348 | 2 (0.6–3.7) | 4 | 310 | 1 (0.3–3.2) |
| <i>Chlamydophila</i> spp. | 63 | 352 | 18 (14.0–22.3) | 87 | 348 | 25 (20.5–30.0) | 49 | 313 | 16 (11.8–20.1) |
| Pestiviruses | 322 | 352 | 91 (88.0–94.1) | 315 | 352 | 90 (80.7–88.5) | 317 | 352 | 90 (86.4–92.9) |
| BRSV | 286 | 325 | 81 (76.7–85.1) | 299 | 352 | 85 (80.7–88.5) | 294 | 352 | 84 (79.2–87.2) |
| <i>M. conjunctivae</i> | 290 | 352 | 82 (78.0–86.2) | 260 | 352 | 74 (69.0–78.4) | 307 | 352 | 87 (83.2–90.5) |

^a Binomial exact one-sided, 97.5% CI.

TABLE 3. Number of animals tested, positive animals, and seroprevalences for selected pathogens in chamois in three valleys of the Province of Bergamo (Italy).

| Pathogen | Val Brembana | | | Val Seriana | | | Val di Scalve | | |
|---------------------------|----------------|--------|-----------------------|----------------|--------|-----------------------|----------------|--------|-----------------------|
| | No. of animals | | Prevalence (%) | No. of animals | | Prevalence (%) | No. of animals | | Prevalence (%) |
| | Positive | Tested | (95% CI) ^a | Positive | Tested | (95% CI) ^a | Positive | Tested | (95% CI) ^a |
| <i>Brucella</i> spp. | 0 | 13 | 0 (0.0–24.7) | 0 | 17 | 0 (0.0–19.5) | 0 | 19 | 0 (0.0–17.6) |
| <i>Toxoplasma gondii</i> | 1 | 24 | 4 (0.1–21.1) | 3 | 32 | 9 (19.8–25.0) | 1 | 50 | 2 (0.0–10.6) |
| <i>Neospora caninum</i> | 4 | 30 | 13 (3.7–30.7) | 6 | 16 | 38 (15.2–64.5) | 4 | 22 | 18 (5.1–40.2) |
| <i>Chlamydophila</i> spp. | 0 | 31 | 0 (0.0–11.2) | 0 | 32 | 0 (0.0–10.8) | 0 | 43 | 0 (0.0–8.2) |
| BHV-1 | 0 | 23 | 0 (0.0–14.8) | 0 | 27 | 0 (0.0–12.7) | 0 | 50 | 0 (0.0–7.1) |
| BPIV-3 | 0 | 13 | 0 (0.0–24.7) | 1 | 27 | 4 (0.0–19.0) | 14 | 50 | 28 (16.2–42.5) |
| Pestiviruses | 0 | 25 | 0 (0.0–13.7) | 0 | 27 | 0 (0.0–12.7) | 18 | 46 | 39 (25.0–54.6) |
| BRSV | 5 | 20 | 25 (8.6–49.1) | 10 | 26 | 38 (20.2–59.4) | 21 | 50 | 42 (28.1–56.8) |
| <i>M. conjunctivae</i> | 34 | 202 | 16 (12.0–22.8) | 2 | 18 | 11 (1.9–36.0) | 4 | 16 | 25 (8.3–52.5) |

^a Binomial exact one-sided, 97.5% CI.

cometti et al., 2002). This is probably due to an IKC outbreak that occurred in the Val Brembana immediately before the collection of our samples.

For chamois, antibody prevalence estimates for pestivirus ($\chi^2=24.9_{(df=2)}$, $P<0.0001$) and BPIV-3 ($\chi^2=10.5_{(df=2)}$; $P=0.005$) differed significantly between the valleys (Table 3). However, we found only moderate values for the contingency coefficients (Cramer $\Phi_{\text{Pestivirus}}=0.54$ and Cramer $\Phi_{\text{PI3}}=0.34$). Antibodies to pestiviruses have been detected in chamois in the western Alps (Olde Riekerink, 2005), and pestivirus infections and related disease have been reported from Pyrenean chamois (Hurtado et al., 2004). The presence of antibodies to BRSV in chamois also may be significant, because this virus may have been associated with epidemic pneumonia in chamois in an area bordering west Val Brembana (Citerio et al., 2003).

Antibodies to *Chlamydophila* spp. were not found in either chamois or roe deer in

this study, but seropositive results have been reported for these species (Cubero-Pablo et al., 2000). In addition, *Chlamydophila abortus* was recently detected by polymerase chain reaction from a mummified fetus from a healthy roe deer from our study area (Gaffuri et al., 2003). Antibodies to BHV-1 also were not detected, and although serologic cross-reactions can occur among alphaherpesviruses (Keuser et al., 2004), we cannot discount the possible presence of other herpesviruses in these populations.

Based on the higher prevalence of antibodies in sheep compared with both roe deer and chamois, it is probable that sheep represent a potentially important source of infection for wild ruminants. However, these high prevalence estimates from sheep may be partially influenced by herd reorganization and the routine mixing of small groups of animals. Our results confirm the absence of brucellosis in this area but demonstrate the potential circulation and transmission of several impor-

tant pathogens between species. To better understand the nature and significance of these interactions, additional clinical data are needed as well as the isolation and molecular characterization of these etiological agents.

The authors are grateful to the Public Veterinary service of the Province of Bergamo, and, in particular, to E. Testa and D. Frosio for collaboration. We also thank T. Ambrosi for contributions as both a veterinarian and a hunter. The hunting districts of the Province of Bergamo and the state gamekeepers also are acknowledged for their cooperation. We thank R. Coates, Centro linguistico dell'Università di Brescia, for linguistic revision. Special thanks are due to C. Garbarino who contributed to the beginning of the study. M.G. was supported by the Fund for Research on Infectious Keratoconjunctivitis, Chur (Switzerland), and P.L. was supported in part by a grant from the University of Milan, Italy (FIRST 2002).

LITERATURE CITED

- ABRAMSON, J. H., AND P. M. GAHLINGER. 1993–2000. Computer programs for epidemiologists (PEPI), version 3.01. Brixton Books, Llanidloes, Wales.
- BELLOY, L., M. GIACOMETTI, E. ABDO, J. NICOLET, M. KRAWINKLER, M. JANOVSKY, U. BRUDERER, AND J. FREY. 2001. Detection of specific *Mycoplasma conjunctivae* antibodies in the sera of sheep with infectious keratoconjunctivitis. *Veterinary Research* 32: 155–164.
- , M. JANOVSKY, E. M. VILEI, P. PILO, M. GIACOMETTI, AND J. FREY. 2003. Molecular epidemiology of *Mycoplasma conjunctivae* in Caprinae: Transmission across species in natural outbreaks. *Applied and Environmental Microbiology* 69: 1913–1919.
- BROCCHI, E., P. CORDIOLI, A. BERLINZANI, D. GAMBA, AND F. DE SIMONE. 1993. Development of a panel of antipestivirus monoclonal antibodies useful for virus identification and antibody assessment. *In Proceedings: 2nd symposium on pestiviruses, European Society for Veterinary Virology, Nancy, France, 1–3 October 1992*, S. Edwards (ed.). Fondation Marcel Mérieux, Lyon, France, pp. 215–218.
- CITTERIO, C. V., C. LUZZAGO, M. SALA, G. SIRONI, P. GATTI, A. GAFFURI, AND P. LANFRANCHI. 2003. Serological study of a population of alpine chamois (*Rupicapra r. rupicapra*) affected by an outbreak of respiratory disease. *The Veterinary Record* 153: 592–596.
- CIUCHINI, F., AND R. FARINA. 1991. Primo progetto di standardizzazione delle tecniche sierologiche per la diagnosi delle brucellosi animali: Sieroagglutinazione rapida con antigene al rosa bengala SAR-Ag: RB, siero-agglutinazione lenta in micrometodo SAL-mi., fissazione del complemento miniaturizzata FdC-mi., Istituto Superiore di Sanità, Roma, Italy, 91/16: 71.
- CONRAD, P. A., K. SVERLOW, M. ANDERSON, J. ROWE, R. BONDURANT, G. TUTER, R. BREITMEYER, C. PALMER, M. THURMOND, A. ARDANS, J. P. DUBEY, G. DUHAMEL, AND B. BARR. 1993. Detection of serum antibody response in cattle with natural or experimental *Neospora* infections. *Journal of Veterinary Diagnostic Investigations* 5: 572–578.
- CONTINI, A., AND P. COSSU. 1975. Presenza di anticorpi inibenti la emoagglutinazione da virus PI-3 in sieri ovini con affezioni respiratorie. *Atti Società Italiana delle Scienze Veterinarie* 29: 686.
- CUBERO-PABLO, M. J., M. PLAZA, L. PÉREZ, M. GONZÁLEZ, AND L. LEÓN-VIZCANO. 2000. Seroepidemiology of chlamydial infections of wild ruminants in Spain. *Journal of Wildlife Diseases* 36: 35–47.
- DE SIMONE, F., E. BROCCHI, Y. L. ARCHETTI, D. GAMBA, AND E. FONI. 1986. Diagnosi diretta e sierologica della malattia da virus respiratorio sinciziale mediante anticorpi monoclonali. *Atti Società Italiana delle Scienze Veterinarie* 40: 903–990.
- D. M. 2 Luglio. 1992. n.453 “Regolamento concernente il piano nazionale per la eradicazione della brucellosi negli allevamenti ovini e caprini” G.U. n.276 del 23 Novembre 1992. Istituto Poligrafico e Zecca dello Stato, Roma, Italy.
- DUBEY, J. P. 1993. *Toxoplasma, Neospora, Sarcocystis* and other tissue cyst-forming coccidian of humans and animals. *In Parasitic protozoa*, Vol. 6. 2nd Edition. J. P. Kreier (ed.). Academic Press, San Diego, California, pp. 1–158.
- FERROGLIO, E., AND L. ROSSI. 2001. Prevalence of *Neospora caninum* antibodies in wild ruminants from the Italian Alps. *Veterinary Record* 148: 754–755.
- , ———, AND S. GENNERO. 2000. Lung-tissue extract as an alternative to serum for surveillance for brucellosis in chamois. *Preventive Veterinary Medicine* 43: 117–122.
- GAFFURI, A., S. MAGNINO, AND L. CAPITANIO. 2003. Rilevamento di *Chlamydophila abortus* in un feto mummificato di capriolo. *In Proceedings: “6° Convegno internazionale malattie infettive nell’arco alpino,” Siusi allo Sciliar BZ. Italy, 20–22 March 2003*, pp. 79–80. Provincia autonoma di Bolzano-Alto Adige, Bolzano, Italy.
- GENNERO, M. S., P. G. MENEGUZ, M. MANDOLA, L. MASOERO, D. DE MENECHI, AND L. ROSSI. 1993.

- Indagini sierologiche su ruminanti selvatici in Piemonte. *Atti Società Italiana delle Scienze Veterinarie* 47: 979–983.
- GIACOMETTI, M., M. JANOVSKY, H. JENNY, J. NICOLET, L. BELLOY, E. GOLDSCHMIDT-CLERMONT, AND J. FREY. 2002. *Mycoplasma conjunctivae* infection is not maintained in Alpine chamois in eastern Switzerland. *Journal of Wildlife Diseases* 38: 297–304.
- GONDIM, L. F., M. M. MCALLISTER, W. C. PITT, AND D. E. ZEMBLICK. 2004. Coyotes (*Canis latrans*) are definitive hosts of *Neospora caninum*. *International Journal for Parasitology* 34: 159–161.
- GRAAT, E. A. M., K. FRANKENA, AND H. BOS. 1997. Principles and methods of sampling in animal disease surveys. In *Application of quantitative methods in veterinary epidemiology*. J. P. T. M. Noordhuizen, K. Frankena, C. M. van der Hoofd, and E. A. M. Graat (eds.). Wageningen Pers, Wageningen, The Netherlands, pp. 31–62.
- HENKEN, A. M., E. A. M. GRAAT, AND J. CASAL. 1997. Measurement of disease frequency. In *Application of quantitative methods in veterinary epidemiology*. J. P. T. M. Noordhuizen, K. Frankena, C. M. Van der Hoofd and E. A. M. Graat (eds.). Wageningen Pers, Wageningen, The Netherlands, pp. 63–97.
- HURTADO, A., G. ADURIZ, N. GOMEZ, B. OPORTO, R. A. JUSTE, S. LAVIN, J. R. LOPEZ-OLVERA, AND I. MARCO. 2004. Molecular identification of a new pestivirus associated with increased mortality in the Pyrenean chamois (*Rupicapra rupicapra pyrenaica*) in Spain. *Journal of Wildlife Diseases* 40: 796–800.
- JEKEL, J. F., J. G. ELMORE, AND D. L. KATZ. 1996. Bivariate Analysis. In *Epidemiology, biostatistics and preventive medicine*. W. B. Saunders, London, UK, 267 pp.
- KEUSER, V., F. SCHYNTS, B. DETRY, A. COLLARD, B. ROBERT, A. VANDERPLASCHEN, P. P. PASTORET, AND E. THIRY. 2004. Improved antigenic methods for differential diagnosis of bovine, caprine, and cervine Alphaherpesviruses related to bovine Herpesvirus 1. *Journal of Clinical Microbiology* 42: 1228–1235.
- MAGNINO, S., P. G. VIGO, M. FABBI, M. COLOMBO, C. BANDI, AND C. GENCHI. 1999. Isolation of a bovine *Neospora* from a newborn calf in Italy. *The Veterinary Record* 144: 456.
- O. I. E. Manual. 1992. Enzootic abortion of ewes (ovine chlamydiosis). (B28): 393–399. Office International des Epizooties, Paris, France.
- OLDE RIEKERINK, R. G. M., A. DOMINICI, H. W. BARKEMA, AND A. J. DE SMIT. 2005. Seroprevalence of pestivirus in four species of alpine wild ungulates in the High Valley of Susa, Italy. *Veterinary Microbiology* 108: 297–303.
- WALLS, K. W., AND J. S. REMINGTON. 1983. Evaluation of a commercial latex agglutination method for toxoplasmosis. *Diagnostic Microbiology and Infectious Diseases* 1: 265–271.

Received for publication 16 December 2004.